

CASE REPORT

Iran J Allergy Asthma Immunol

April 2020; 19(2):200-202.

Doi: 10.18502/ijaai.v19i2.2773

Anaphylaxis to Oatmeal and Psocid Crisps

Delara Babaie^{1,2}, and Peter Vadas¹

¹ Division of Clinical Immunology and Allergy, Department of Medicine, St. Michael's Hospital, University of Toronto, Toronto, Canada

² Department of Allergy and Clinical Immunology, Mofid Children's Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Received: 21 May 2019; Received in revised form: 2 November 2019; Accepted: 6 November 2019

ABSTRACT

Occasionally, a seemingly straightforward history of food-induced anaphylaxis may prove to be misleading. Both patients and their physicians have a tendency to attribute the cause of an allergic reaction to the most conspicuous ingredient that had been ingested while overlooking less likely causes. Here, we describe a patient whose history pointed to oatmeal allergy, but skin prick tests to oats and serologic testing for oat-specific IgE were negative. Ultimately, we found that the oatmeal had been contaminated with an allergenic insect, Psocid of the order Psocoptera.

Keyword: Food allergy; Oatmeal allergy; Psocid

INTRODUCTION

The prevalence of self-reported food allergies is very much higher than the prevalence of food allergies confirmed by a detailed clinical history, utilization and interpretation of appropriate tests.^{1,2} Modalities recommended in the 2010 Expert Panel Guidelines include a detailed medical history and physical examination, skin prick tests (SPTs), allergen-specific IgE (sIgE) measurements, and oral food challenge (OFC).³ Sometimes identifying the causative allergen is diagnostically challenging, especially when the evaluations are inconclusive or indeterminate. In addition, a mistaken diagnosis may lead to unnecessary avoidance of food, a false sense of security and the potential for recurrent reactions.

Here, we describe a patient suspected to have an oatmeal allergy, but investigations for oat allergy were negative. Eventually, we found that her homemade oatmeal crisps had been contaminated by an allergenic insect, Psocid of the order *Psocoptera*, and it could explain all her symptoms.

CASE REPORTS

A 61-year-old female with allergic rhinitis and hypertension was referred to the Anaphylaxis Clinic at St. Michael's Hospital, a tertiary care teaching hospital fully affiliated with the Faculty of Medicine, University of Toronto. Several months earlier, she had been preparing an oatmeal crisp, mixing packaged oatmeal with butter and brown sugar with her bare hands. Within 5 minutes, she developed itchy palms but no other cutaneous or visceral manifestations. She later ingested a small portion of the baked crisp and felt "heaviness in her stomach". Several weeks later, she

Corresponding Author: Peter Vadas, MD, PhD;
Division of Clinical Immunology and Allergy, Department of
Medicine, St. Michael's Hospital, University of Toronto, Toronto,
Canada. Tel: (+416) 8645 777, E-mail: vadasp@smh.ca

Anaphylaxis to Psocide

again prepared oatmeal cookies with bare hands but did not have any immediate symptoms on contact. While cleaning up, she believes that she inhaled some airborne oatmeal dust. Several minutes later, she developed paroxysms of sneezing, nasal congestion, itchy palms, and soles, as well as itchy, red, watery eyes. She went to an emergency department where she became lightheaded and vomited. She was treated with IV diphenhydramine and prednisone and discharged 6 hours later in stable condition. Prior to these reactions, she had eaten oatmeal on a number of occasions as a part of Harvest Crunch oatmeal cereal without any adverse symptoms, suggesting a negative oral food challenge. She has avoided eating oats subsequent to the reactions above.

She was a retired dental hygienist. Her physical examination was unremarkable. SPT to food allergens were all non-reactive using commercial diagnostic extracts, including skin tests to oat, wheat, rye, barley, corn, and rice. SPT to fresh oatmeal was similarly non-reactive. ImmunoCAP testing for sIgE to oat was <0.01 KIU/ml (<0.35 KIU/mL is negative). A formal graded oral challenge with an oatmeal cookie was planned for her. In the meantime, she mistakenly ate part of the oatmeal cookie at a party and had no untoward symptoms.

When she was preparing some oatmeal crisps in anticipation of a graded oral challenge, she noted numerous foreign bodies in her oatmeal. A sample was submitted to the Canadian Food Inspection Agency, whose investigations showed a heavy infestation of adult and nymph forms of psocids.

DISCUSSION

Psocids, commonly known as booklice, belong to the order *Psocoptera*. They are tiny, soft-bodied, winged insects, with worldwide distribution.⁴ Their biology and ecology differ among various species. They range in size from 1–10 mm in length.⁵ Some species of Psocids, such as *Liposcelisbostrychophila*, commonly infest stored products, such as oats. Psocids are able to spread fungal pathogens, which can threaten human health.⁶ These insects have been reported to cause allergic reactions in sensitized individuals,⁷ as can mealworm infestations in stored grain⁸ and dust mite infestation in pancake batter.⁹

Herein, we describe a patient with an IgE-mediated food allergy, who experienced a contact reaction after

handling dough made from contaminated oatmeal and a subsequent multisystem reaction after inhalation of contaminated oatmeal dust. *In vivo* and *in vitro* testing for oatmeal allergy were negative. She ate oatmeal cookies accidentally in a party without any symptoms. Visual inspection of the oatmeal revealed the presence of an insect contaminant, subsequently identified by the Canadian Food Inspection Agency as a heavy infestation by Psocidlarvae and adults, which can clinically be suggested as an allergenic trigger of her IgE-mediated multisystem reaction.

Recent studies have shown Psocids are more tolerant to some of the insecticides compared to other stored-product insect pests,¹⁰ leading to an increased prevalence in the commercial food supply. *L. bostrychophila* is reported to be an important trigger of respiratory allergy,¹⁰ and causes IgE mediated responses in up to 20% of the allergic population studied.¹¹ A novel 26 kDIgE-binding protein from *PsocidLip b 1* has been found^{10,11} which does not share sequence homology with any previously described allergenic proteins.¹² Moreover, Several allergenic proteins have been characterized in storage mite-infested foods, including TP *Lepidoglyphus destructor* (LD) and *Tyrophagusputrescentiae* (TP) which also cause allergy in human.¹³

Allergic reactions are not necessarily due to the most conspicuous ingredient amongst the foods ingested. A formal investigation is needed to identify the cause so that appropriate avoidance measures may be implemented. In this case, we report the presence of Psocids contaminating oatmeal, clinically could trigger of this patient's multisystem allergic reaction, indeed, detection of *PsocidLip b 1 protein* is necessary to confirm the diagnosis.

Conflict of Interest

The authors have no conflict of interest regarding this case report. The patient signed the informed consent to report the case.

ACKNOWLEDGEMENTS

The authors would like to thank the Canadian Food Inspection Agency.

REFERENCES

1. Soller L, Ben-Shoshan M, Harrington DW, Fragapane J,

- et al. Overall prevalence of self-reported food allergy in Canada. Vol. 130, The Journal of allergy and clinical immunology. United States; 2012. p. 986–8.
2. Warren CM, Gupta RS, Smith BM, Davis MM, Jiang J, Blumenstock JA, et al. Prevalence and Severity of Food Allergies Among US Adults. *JAMA Netw Open*. 2019;2(1):e185630.
 3. Boyce JA, Assa'ad A, Burks AW, Jones SM, Sampson HA, Wood RA, et al. Guidelines for the diagnosis and management of food allergy in the United States: report of the NIAID-sponsored expert panel. *J Allergy Clin Immunol*. 2010 Dec;126(6 Suppl):S1-58.
 4. Phillips TW, Throne JE. Biorational approaches to managing stored-product insects. *Annu Rev Entomol*. 2010;55:375–97.
 5. Athanassiou CG, Arthur FH, Opit GP, Throne JE. Insecticidal effect of diatomaceous earth against three species of stored-product psocids on maize, rice, and wheat. *J Econ Entomol*. 2009 Aug;102(4):1673–80.
 6. Obr S. Psocoptera of food-processing plants and storages, dwellings and collections of natural objects in Czechoslovakia. *Acta Entomol Bohemoslov*. 1978;75:226–42.
 7. Turner BD. *poscelis bostrychophila* (Psocoptera: Liposcelididae), a stored-product pest in the UK. *Int J Pest Manag*. 1994;40:179–90.
 8. Schroeckenstein DC, Meier-Davis S BR. Occupational sensitivity to *Tenebrio molitor* Linnaeus (yellow mealworm). *Allergy Clin Immunol*. 1990;86(2):182–8.
 9. Masaki K, Fukunaga K, Kawakami Y, Haque R. Rare presentation of anaphylaxis: pancake syndrome. *BMJ Case Reports CP* [Internet]. 2019;12(3). Available from: <https://casereports.bmj.com/content/12/3/e228854>
 10. Ishibashi O, Sakuragi K, Fukutomi Y, Kawakami Y, Kamata Y, Sakurai M, et al. Lip b 1 is a novel allergenic protein isolated from the booklouse, *Liposcelis bostrychophila*. *Allergy*. 2017 Jun;72(6):918–26.
 11. Patil MP, Niphadkar P V, Bapat MM. Psocoptera spp. (book louse): a new major household allergen in Mumbai. *Ann Allergy Asthma Immunol*. 2001 Aug;87(2):151–5.
 12. Fukutomi Y, Kawakami Y, Taniguchi M, Saito A, Fukuda A, Yasueda H, et al. Allergenicity and cross-reactivity of booklice (*Liposcelis bostrychophila*): a common household insect pest in Japan. *Int Arch Allergy Immunol*. 2012;157(4):339–48.
 13. Jakubas-Zawalska J, Asman M SK. Sensitization to the storage mites *Lepidoglyphus destructor* and *Tyrophagus putrescentiae* (Acari, Sarcoptiformes, Astigmatina) in a suburban population in Southern Poland. *Ann Parasitol*. 2017;63(3):183–8.