

ORIGINAL ARTICLE

Iran J Allergy Asthma Immunol

August 2020; 19(4):386-396.

Doi: 10.18502/ijaai.v19i4.4113

The Relaxant Effect of *Plantago Major* on Rat Tracheal Smooth Muscles and Its Possible Mechanisms

Javad Boskabadi^{1,2}, Saeideh Saadat³, and Mohammad Hossein Boskabady^{1,4}

¹ Neurogenic Inflammation Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

² Department of Clinical Pharmacy, School of Pharmacy, Mazandaran University of Medical Sciences, Sari, Iran

³ Department of Physiology, School of Medicine, Zahedan University of Medical Sciences, Zahedan, Iran

⁴ Department of Physiology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

Received: 6 December 2019; Received in revised form: 13 March 2020; Accepted: 20 March 2020

ABSTRACT

This study was conducted to evaluate the possible mechanisms of the relaxant effects of hydroalcoholic extract of *Plantago major* (*P. major*) on tracheal smooth muscle (TSM) in rats.

The effects of cumulative concentrations of *P. major* (5, 10, 20 and 40 mg/mL) and theophylline (0.2, 0.4, 0.6 and 0.8 mM) were evaluated on pre-contracted TSM with 10 μ M methacholine or 60 mM KCl. To determine the possible mechanisms, the relaxant effect of the plant was also examined on incubated TSM with atropine, indomethacin, chlorpheniramine, glibenclamide, diltiazem, papaverine, and propranolol.

The results indicated concentration-dependent relaxant effects for *P. major* in non-incubated TSM contracted by methacholine or KCl. There was no statistically significant difference in the relaxant effects of *P. major* between non-incubated and incubated tissues with indomethacin, papaverine, and propranolol. However, the relaxant effects of *P. major* in incubated tissues with atropine ($p < 0.01$ to $p < 0.001$), chlorpheniramine ($p < 0.05$ to $p < 0.001$), glibenclamide ($p < 0.05$), or diltiazem ($p < 0.01$) were significantly lower than non-incubated TSM.

P. major indicated relatively potent relaxant effects which were lower than those of theophylline. Muscarinic and histamine (H₁) receptors inhibition, as well as calcium channel blocking and potassium channel opening effects are suggested to contribute to the TSM relaxant effect of the plant.

Keywords: Histamine (H₁) receptor; Muscarinic receptor; *Plantago major*; Smooth muscle; Trachea

INTRODUCTION

Plantago major L. (*P. major*), broadleaf plantain, is an amazing herb that belongs to the Plantaginaceae family.¹ The seeds of the *P. major* are ovate shape and small size (0.4–0.8_0.8–1.5 mm) with a bit bitter taste.²

It grows in compressed soils and is plentiful beside routes, roadsides, and other areas. It propagates by seeds, which are held on the long, narrow spikes that rise well above the foliage.³ This plant grows in regions that weed is common and widely spread and naturalized in

Corresponding Author: Mohammad Hossein Boskabady, MD, PhD; Neurogenic Inflammation Research Center, Mashhad University of

Medical Sciences, Mashhad, Iran. Tel: (+98 51) 3882 8565, Fax: (+98 51) 3882 8564, E-mail: boskabady@ums.ac.ir

the world. *P. major* is native to northern and central Asia and most of Europe. In 4000 years ago, *P. major* was spread by the man from Europe throughout the world.⁴

P. major is a medicinal plant that used for the treatment of various conditions (e.g., skin problems, respiratory diseases, digestive tracts, and urinary infection).² A lot of beneficial effects have been reported for *P. major* leaves extracts including anti-inflammation, antitumor, antibacterial, antifungal⁵, antiviral, analgesic, antioxidant, anti-ulcer, renal and hepatoprotective⁶, and immune-modulating activities.⁷⁻⁹ Chemical studies on various *P. major* extracts indicated that phenol groups are major compounds of the plant. Terpenoids, flavonoids as organic acids, alkaloids and glycosides are observed in *P. major* extracts (Figure 1).¹⁰ Also, Ursolic acid (0.22%) and oleanolic acid (0.07%) are terpenoids compounds of *P. Major*.^{10,11}

The above-mentioned compounds found in almost all parts of the plant. Bioactivity of *P. major* leaves contained the secondary metabolites is attributed to these chemical constituents.¹² It has been demonstrated that the anti-inflammatory effect of the plant may relate to ursolic acid. This terpenoid compound is a selective inhibitor of cyclooxygenase-2 (COX-2), that responsible for prostaglandins biosynthesis; so, *P. major* have an anti-inflammatory effect by COX-2 inhibition.¹²




Figure 1. Some chemical structural of *Plantago major* (*P. Major*) compounds including baicalein (flavonoid), aucubin (irioid glycoside), oleanolic acid (terpenoid) and plantagonin (alkaloid)

Leaves of the plant are rich sources of essential fatty acids (18:2 ω and 18:3 ω) and also of carotenes.¹³ *P. major*, also contain some other effective components including ferulic acid, ursolic acid, baicalein, apigenin, benzoic acid, chlorogenic acid, ascorbic acid, citric acid, oleanolic acid, salicylic acid.¹⁴ It is believed that the most important chemicals ingredients in *P. major* are baicalein (a flavonoid) and aucubin (an irioid glycoside) which are biologically active agents (Figure 1).²

In recent years, in both non-developed and developed countries, chemical synthetic drugs have been extensively substituted with herbal medicines.¹⁵ This plant can relax smooth muscles for example, in clinical practice, many brands of *P. major* are used to decrease gastrointestinal movements and diarrhea management.¹⁰ On the other hand, the effect of *P. major* on respiratory and allergic diseases was shown.¹⁶

Based on previous studies, the present study was designed to assess the possible mechanisms responsible for the relaxant effect of hydroalcoholic extract of *P. major* on rat tracheal smooth muscle (TSM). Additionally, to evaluate the effect of *P. major* on β_2 -adrenergic, muscarinic and histamine (H_1) receptors, prostacyclin and COX-2 pathway, potassium and calcium channels, as well as phosphodiesterase activity, the relaxant effect of *P. major* was examined on TSM incubated with propranolol, atropine, chlorpheniramine, glibenclamide, diltiazem, indomethacin, and papaverine respectively.

P. major indicated concentration-dependent relaxant effects in non-incubated TSM contracted by potassium chloride (KCl) and methacholine which was reduced in incubated tissue with atropine chlorpheniramine, glibenclamide, and diltiazem. The results indicated a potent relaxant effect of *P. major* which suggested to be due to muscarinic and histamine (H_1) receptors inhibition, calcium channel blocking, and potassium channel opening effects.

MATERIALS AND METHODS

Materials

P. major seeds were obtained from a grocery market in Mashhad, Khorasan Razavi province, Iran, October 2017. This plant was recognized and identified by Dr. Iranshahi (PharmD, Ph.D. in Pharmacognosy), a professor in the Department of Pharmacognosy, Faculty of Pharmacy, Mashhad University of Medical

Sciences, Mashhad, Iran.

KCl was obtained from Merck (Darmstadt, Germany). Methacholine, atropine, chlorpheniramine, indomethacin, diltiazem, glibenclamide, propranolol and papaverine were also purchased from Sigma Chemical Co. Ltd (UK).

Preparation of *P. Major* Extract

P. major seeds were carefully dried and cleaned and powdered with mixer. Next, 100 g of powdered seeds were poured into Erlenmeyer flask and soaked up with a hydro-ethanolic solution (70%) for 3 days, with stirring and shaking at 37°C (in shaker-stirrer).

To remove alcohol, the mixture was then filtered by Eyela (Heidolph, Germany) rotary evaporator and the resultant solution was concentrated under reduced pressure at 45°C for 24 h in an oven (for drying). The yield extract was 13% and it was stored at -70°C until use.

Animals

The experiments were performed on fifty-one male Wistar rats (7 weeks, weighting 200-250 g). The animals were purchased from the Animal House, Faculty of Medicine, Mashhad University of Medical Sciences (Mashhad, Iran) and maintained under controlled conditions with 12 h/ 12 h light/dark cycle at 22±2 °C. Water and food were always accessible *ad libitum* to animals. This study was approved by the Committee on Animal Research of the Mashhad University of Medical Sciences for Animal Experiments (Approval No. 930658).

Tissue Preparation

After sacrificing the animals, the chest was opened and segmental resection of the trachea was prepared including four rings. The tissues mounted in a 10 mL

organ bath containing Krebs- Henseleitsolution (KHS), and maintained at 37±0.5 °C with isometric tension of 1 g as previously described¹⁷. The isometric transducer (MLT0202, AD Instruments, Australia) linked to a power lab system (Power Lab 8/30, ML870, AD Instruments, Australia) was used to measure contraction and relaxation responses in all experiments.

Experimental Design

In various groups, the relaxant effect of *P. major* extract was measured according to previous studies^{18,19} on incubated or non-incubated tissues with different drugs for 10 minutes as described in Table 1.

To examine the relaxant effect of *P. major*, rat TSM was contracted by methacholine for 7 minutes or KCl for 5 minutes in the non-incubated (n=10) and incubated tissues with 1 µM atropine (n=11), 1 µM indomethacin (n= 8), 1 µM chlorpheniramine (n=8), 5 µM diltiazem (n=8), 1 µM glibenclamide (n=8), 1 µM propranolol (n= 8) or 50 µM papaverine (n=8).

Cumulative concentrations of *P. major* extract (5, 10, 20. and 40 mg/mL) were added on pre-contracted TSM with methacholine or KCl every 5 minutes and at the end of the intervals, the relaxation was recorded in each experiment. Concentrations of *P. major* were chosen based on previous studies.¹⁸ The percent relaxation of each concentration of *P. major* or theophylline was plotted relative to the maximum contraction due to methacholine or KCl to make a concentration-response graph. Theophylline (0.2, 0.4, 0.6, and 0.8 mM) and saline (1 mL) were also examined as positive control and negative control respectively. Furthermore, the effective concentration of *P. major* causing 50% of maximum response (EC₅₀) was also measured from concentration-response curves.¹⁹

Table 1. The protocol of the study and the methods of evaluating various mechanisms of the relaxant of the effect of *Plantago major* (*P. major*) on tracheal smooth muscle (TSM)

TSM Contracting agent	Condition	Incubating substance	Mechanisms
60 mM KCl	Non-incubated tissues (n=10)		
	Incubated tissues	1 µM atropine (n=11) 1 µM indomethacin (n=8)	Muscarinic receptor inhibition Cyclooxygenase inhibition
10 µM methacholine	Non-incubated tissues (n=10)		
	Incubated tissues	1 µM chlorpheniramine (n=8)	Histamine (H ₁) receptor inhibition
		5 µM diltiazem (n=8)	Calcium channel blocking
		1 µM glibenclamide (n=8)	Potassium channel opening
		1 µM propranolol (n=8)	β-adrenoceptor stimulation
	50 µM papaverine (n=8)	Phosphodiesterase inhibition	

Statistical Analysis

All data were presented as mean±SEM. Statistical comparisons of the results were analyzed using a one-way analysis of variance (ANOVA) followed by Tukey's Multiple Comparison Test. Statistically significant was considered as $p<0.05$. The correlation between the relaxant effect of *P. major* and theophylline with their concentrations was assessed by Pearson's correlation coefficient.

RESULTS

P. major Induced Relaxant Effect in non-incubated TSM Contracted by KCl or Methacholine

Concentration-dependent and significant relaxant effects were exhibited for *P. major* in non-incubated TSM contracted by KCl ($p<0.01$ to $p<0.001$ for the last three higher concentrations of the extract). However, the relaxant effect of all concentrations of the *P. major* extract, were significantly lower than those of theophylline ($p<0.001$ for all concentrations, Figure 2a).

P. major extract demonstrated significant and concentration-dependent relaxant effects in non-incubated TSM contracted by methacholine ($p<0.001$ for three last concentrations, Figure 2b). In methacholine-induced contraction also, the relaxant effects of all concentrations of the extract were significantly lower than those of theophylline ($p<0.01$ for its high concentration and $p<0.001$ for other concentrations, Figure 2b).

P. major Induced Relaxant Effect in Incubated TSMs Contracted by KCl

In tissues incubated with atropine, the relaxant effects of three first concentrations of *P. major* were significantly lower than non-incubated TSM ($p<0.01$ to $p<0.001$, Figure 3a). However, in incubated tissues with indomethacin, the relaxant effect of a high concentration of *P. major* was significantly higher than the non-incubated condition ($p<0.05$, Figure 3b).

No statistically significant difference was observed among EC_{50} values of the extract in non-incubated TSM contracted by KCl (11.06 ± 3.87) or tissues incubated with atropine (18.75 ± 5.63) and indomethacin (10.45 ± 5.62 , Figure 3c).




Figure 2. Concentration-response relaxant effect of hydroalcoholic extract of *P. major* (5, 10, 20, 40 mg/mL), (n=10) and theophylline (0.2, 0.4, 0.6, 0.8 mM) (n=6) on (a) KCl (60 mM) or (b) methacholine (10 μ M) induced contraction of tracheal smooth muscle (TSM) in non-incubated tissues. Data are shown as mean \pm SEM.***: $p<0.001$ compared to saline (NS). +++: $p<0.001$ and ++: $p<0.01$ compared to the effect of theophylline.

P. major Induced Relaxant Effect in Incubated TSMs Contracted by Methacholine

The relaxant effects of three last concentrations of *P. major* in incubated tissues with chlorpheniramine and the relaxant effect of its second concentration (10 mg/mL) in incubated tissues with glibenclamide or diltiazem were significantly lower compared to non-incubated TSM ($p<0.05$ to $p<0.001$, Figure 4a-c).




Figure 3. The concentration-response relaxant effect of hydroalcoholic extract of *Plantago major* (*P. major*) on KCl (60 mM) induced contraction of tracheal smooth muscle (TSM) in non-incubated (n=10) and incubated tissues with (a) atropine (1 μ M, n=11) and (b) indomethacin (1 μ M, n=8). (c) EC₅₀ values of hydroalcoholic extract of *P. major* induced relaxation obtained on contracted TSMs of the rat with KCl in non-incubated and incubated tissues with atropine and indomethacin. Data are shown as mean \pm SEM. ***: $p < 0.001$, **: $p < 0.01$ and *: $p < 0.05$ compared to saline (NS). +: $p < 0.05$ compared to non-incubated tissues. Statistical Analyses were performed using one-way analysis of variance (ANOVA) followed by Tukey's multiple comparison tests

However, the relaxant effects of various concentrations of *P. major* in incubated tissues with papaverine or propranolol were not statistically different from non-incubated tissues (Figures 4d and 4e).

There was no statistically significant difference among EC₅₀ values of the extract of the plant in non-incubated tissue contracted by methacholine (13.55 \pm 6.25) and TSM incubated with chlorpheniramine (16.90 \pm 6.20), glibenclamide (16.67 \pm 6.62), diltiazem (14.70 \pm 5.10), papaverine (14.17 \pm 3.47) or propranolol (16.05 \pm 6.64), (Figure 4f).

Also, there were no statistically significant differences in the relaxant effect of various concentrations of *P. major* obtained in KCl induced contraction with those in TSM contracted by methacholine (Figure 5).

Correlations of the Relaxant Effect of *P. major* or Theophylline with Their Concentrations

In non-incubated TSM contracted by KCl or methacholine, there were significant correlations between the relaxant effect of theophylline ($p < 0.001$ and $p < 0.01$, respectively) or *P. major* ($p < 0.01$ and $p < 0.001$, respectively) and their concentrations (Table 2).

Significant correlations were also seen between the relaxant effect of *P. major* and its concentrations in KCl or methacholine induced contraction of TSM, incubated with atropine ($r = 0.728$, $p < 0.001$), indomethacin ($r = 0.785$, $p < 0.001$), chlorpheniramine ($r = 0.721$, $p < 0.01$), diltiazem ($r = 0.750$, $p < 0.001$), glibenclamide ($r = 0.703$, $p < 0.01$), papaverine ($r = 0.910$, $p < 0.001$), or propranolol ($r = 0.828$, $p < 0.001$, Table 2).

Relaxant Effect of *Plantago Major*




Figure 4. The concentration-response relaxant effect of hydroalcoholic extract of *Plantago major* (*P. major*) on methacholine (10 μ M) induced contraction of tracheal smooth muscle (TSM) in non-incubated (n=10) and incubated tissues with (a) chlorpheniramine (1 μ M, n=8), (b) glibenclamide (1 μ M, n=8), (c) diltiazem (5 μ M, n=8), (d) papaverine (50 μ M, n=8) and (e) propranolol (1 μ M, n=8). (f) EC₅₀ values of hydroalcoholic extract of *P. major* induced relaxation obtained on contracted rat TSM with methacholine in non-incubated and incubated tissues with chlorpheniramine (n=8), glibenclamide (n=8), diltiazem (n=8), propranolol (n=8) and papaverine (n=8). ***: $p < 0.001$, **: $p < 0.01$ and *: $p < 0.05$ compared to saline (NS). +++: $p < 0.001$, ++: $p < 0.01$ and +: $p < 0.05$ compared to non-incubated tissues. Statistical analyses were performed using a one-way analysis of variance (ANOVA) followed by Tukey's multiple comparison test.




Figure 5. The concentration-response relaxant effect of hydroalcoholic extract of *Plantago major* (*P. major*) on methacholine (10 µM) and KCl (60 mM) induced contraction of non-incubated tracheal smooth muscle (TSM) (n=10). Data are shown as mean ± SEM. ***: $p < 0.001$ and *: $p < 0.05$ compared to saline (NS). Statistical analyses were performed using a one-way analysis of variance (ANOVA) followed by Tukey's multiple comparison test.

Table 2. Relationship between the relaxant effect of *Plantago major* (*P. major*) and theophylline with their concentrations in different experimental groups

Contractile agents	Studied agents	Conditions	R	p value
KCl	<i>P. major</i>	Non-Inc	0.601	$p < 0.01$
		Atropine-Inc	0.728	$p < 0.001$
		Indomethacin-Inc	0.785	$p < 0.001$
Methacholine	Theophylline	Non-Inc	0.790	$p < 0.001$
		Non-Inc	0.873	$p < 0.001$
	<i>P. major</i>	Chlorpheniramine-Inc	0.721	$p < 0.01$
		Diltiazem-Inc	0.750	$p < 0.001$
		Glibenclamide-Inc	0.703	$p < 0.01$
		Papaverine-Inc	0.910	$p < 0.001$
		Propranolol-Inc	0.828	$p < 0.001$
Theophylline	Non-Inc	0.607	$p < 0.01$	

Data were presented as mean±SEM. The correlation between the relaxant effect of *Plantago major* (*P. Major*) and theophylline with their concentrations was assessed by Pearson's correlation coefficient.

DISCUSSION

This study examined the relaxant effect of *P. major* on pre-contracted TSM by methacholine or KCl, and indicated a relatively potent relaxant effect for *P. major*.

The relaxant effects of the medicinal plants on TSM might be produced by various mechanisms such as β_2 -adrenergic receptors stimulation,²⁰ histamine (H_1) receptors inhibition,²¹ calcium channel-blocking,²²

potassium channel opening,²³ muscarinic receptors inhibition,²⁴ methylxanthine activity,²⁵ inhibition of nitric oxide synthase (NOS),²⁶ inhibition of the enzyme phosphodiesterase,²⁷ inhibition of cyclooxygenase (COX) pathway,²⁸ and inhibiting the ATP-sensitive potassium channels.²⁹ The previous study has shown that Flavonoid compounds of *P. major* have a relaxant effect on smooth muscle via potassium channels opening and beta2-adrenergic receptor activation and histamine (H_1) receptors inhibition.¹⁷

Relaxant Effect of *Plantago Major*

In the present study, to evaluate the effect of *P. major* on muscarinic receptors, histamine (H₁) receptors, prostacyclin, and COX-2 inhibitory mechanism, potassium channel opening, calcium channel-blocking, β₂-adrenergic receptors and phosphodiesterase activity, the relaxant effect of *P. major* was examined in TSM incubated with atropine, chlorpheniramine, indomethacin, glibenclamide, diltiazem, propranolol, or papaverine respectively.

In incubated TSM with atropine and contracted with KCl, the relaxant effect of hydroalcoholic extract of *P. major* was examined to assess the contribution of muscarinic receptor inhibitory effect on the relaxant property of the plant. The results of this experiment indicated significantly lower relaxant effects of three first concentrations of *P. major* in incubated tissues with atropine compared to non-incubated TSM EC₅₀ value obtained in TSM incubated with atropine was not statistically different compared to non-incubated

conditions. These results indicated the inhibitory effect of *P. major* on muscarinic receptors which could be contributed in its relaxant effect of the extract on TSM.

The main muscarinic receptor in the smooth muscle is the M₃ receptor which activates phospholipase C and inositol trisphosphate to release calcium from intracellular stores.³⁰ This expresses that a combination of muscarinic receptor antagonists and β₂-adrenoceptor agonists which used in obstructive respiratory diseases, have synergistic effects on airway caliber, so the dose of these drugs in comparison with monotherapy is reduced.³¹ Muscarinic receptors also can induce the production of inflammatory mediators and pro-inflammatory cytokines³² especially in cigarette-smokers and allergic conditions.³³ Lower relaxant effect of all concentrations of *P. major* on incubated tissues with atropine compared to the effects obtained in TSM incubated with other agents also support this mechanism of action for *P. major* (Table 3).

Table 3. Comparison of the relaxant effect *Plantago major* (*P. major*) (percentage change in proportion to the maximum contraction) in different incubated tracheal smooth muscles (TSMs) contracted by 10 μM methacholine (n=8) or 60 mM KCl (n=10)

Incubating substance	Concentration (mg/mL)			
	5	10	20	40
Atropine	0.24 ± 4.11	1.35 ± 4.28	17.34 ± 6.40	31.17 ± 6.73
Indomethacin	33.55 ± 7.98 **	52.43 ± 9.82 **	80.17 ± 13.06 **	92.49 ± 9.52 ***
Chlorpheniramine	11.06 ± 6.88	13.29 ± 3.22	21.27 ± 3.10	41.48 ± 7.29
Diltiazem	6.34 ± 2.53	12.17 ± 3.59	31.33 ± 6.50	48.63 ± 11.58
Glibenclamide	9.35 ± 3.42	17.04 ± 5.72	33.64 ± 10.77	54.39 ± 11.09
Papaverine	4.62 ± 2.61	16.94 ± 7.04	28.23 ± 10.38	62.73 ± 8.43
Propranolol	2.52 ± 8.09	20.11 ± 10.05	58.81 ± 12.45 #	85.81 ± 6.75 ## † ^

Data were presented as mean ± SEM. ***:p<0.001 and **:p<0.01 compared to incubated tissues with atropine. ##:p<0.01 and #:p<0.05 compared to incubated tissues with chlorpheniramine. †:p<0.05 compared to incubated tissues with diltiazem. ^:p<0.05 compared to incubated tissues with glibenclamide.

To study the effect of *P. major* on histamine (H₁) receptor and the involvement of this mechanism in the relaxant effect of the plant, its relaxant effect was examined in TSM incubated with chlorpheniramine and contracted with methacholine. The results of this group expressed significantly lower relaxant effects of three higher concentrations of *P. major* in incubated tissues with chlorpheniramine compared to the non-incubated TSM. Histamine receptors are involved in cell contraction. When these receptors (H₁) activated,

intracellular calcium levels were increased via phospholipase C cascade. On the other hand, histamine receptors have an inflammatory effect by inducing the release of cytokines that can activate Th1 and Th2 immune cells.³⁴ The results indicated the inhibitory (non-competitive) effect of the extract on histamine receptors which could be contributed in its relaxant effect on TSM. In a clinical trial also, *P. major* indicated antihistamine effect in urticarial, which support our finding.³⁵

To evaluate the involvement of prostacyclin mechanism and COX pathways in *P. major* relaxation, its effect was examined in incubated tissues with indomethacin as nonselective COX inhibitor. As shown in the previous study, some compounds of *P. major* extract such as α -linolenic acid, triterpenic acid, ursolic acid, and oleanolic acid have COX inhibitory effect.³⁶ The anti-inflammatory effect of *P. major* was also seen previously⁶ which could be due to its cyclooxygenase inhibitory effect. The significant lower relaxant effect of the highest concentration of the extraction incubated tissues with indomethacin compared to non-incubated condition suggested that the COX inhibitory effect of *P. major* may contribute to its TSM relaxation with a non-competitive manner.

To examine the involvement of calcium channel blocking of *P. major*, the relaxant effect of the extract was studied in incubated TSM with diltiazem. The significant lower relaxant effects of the second concentration of the extract (10 mg/mL) were seen in incubated tissue with diltiazem compared to non-incubated tissues. Therefore, the calcium channel-blocking mechanism may also contribute to the relaxant effect of *P. major* on TSM.

In incubated TSM with glibenclamide, the relaxant effect of *P. major* was evaluated to determine the involvement of potassium channels in the relaxant effect of the plant. The significant lower relaxant effect of the second concentration of the extract (10 mg/mL) was observed. Potassium is an intracellular ion, therefore the opening of potassium channels, leads to cell hyperpolarization and potassium diffusion out of the cell. It is indicated that *P. major* causes potassium diffusion outside the cell³⁷ which suggests the other possible mechanism responsible for the relaxant effect of the plant could be the potassium channel-opening effect.

However, the relaxant effect of the extract in incubated TSM with propranolol and papaverine was not significantly different from those on non-incubated TSM. These results indicated that β 2-adrenergic stimulation and phosphodiesterase inhibitory mechanisms are not contributed to the relaxant effect of *P. major*.

One of the main limitations of this study is the lack of the characterization of the extract of the plant by HPLC which should be performed in a further study. In addition, the other possible mechanisms responsible for the relaxant effect of the plant including its effect on

the non-adrenergic non-cholinergic nervous system should be examined in future studies. The clinical studies for evaluating the bronchodilatory effect of *P. major* in patients with obstructive pulmonary diseases should be also undertaken in the future.

The present study provided novel information about the TSM relaxant effect of *P. major*. Inhibition of muscarinic and histamine (H₁) receptors, calcium channel blocking, and potassium channel opening effects were suggested as possible mechanisms for the relaxant effect of *P. major*.

Taken together, hydroalcoholic extract of *P. major* indicated a relatively potent relaxant effect on TSM which was lower than the effect of theophylline at studied concentrations. Based on the results of this study and the pathophysiology of pulmonary disease, *P. major* can be used as an effective medicinal plant in the treatment of chronic inflammatory respiratory and obstructive pulmonary problems via expressed mechanisms. It is suggested that the efficacy of this plant be studied in future animal studies and clinical trials.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

ACKNOWLEDGEMENTS

This study was financially supported by a grant from the Research Council of Mashhad University of Medical Sciences.

REFERENCES

1. Lukova PK, Karcheva-Bahchevanska DP, Bivolarski VP, Mladenov RD, Iliev IN, Nikolova MM. Enzymatic hydrolysis of water extractable polysaccharides from leaves of *Plantago major* L. *Folia Medica*. 2017;59(2):210-6.
2. Samuelsen AB. The traditional uses, chemical constituents and biological activities of *Plantago major* L. A review. *J Ethnopharmacol*. 2000;71(1):1-21.
3. Cavers PB, Bassett IJ, Crompton CW. The biology of Canadian weeds.: 47. *Plantago lanceolata* L. *Can J Plant Sci*. 1980;60(4):1269-82.
4. Nazarizadeh A, Mikaili P, Moloudizargari M, Aghajanshakeri S, Javaherypour S. Therapeutic uses and pharmacological properties of *Plantago major* L. and its

- active constituents. *J Basic Appl Sci Res.* 2013;3(9):212-21.
5. Abd-Razik B, Hasan H, Murtadha M. The study of antibacterial activity of *Plantago major* and *ceratonia siliqua*. *Iraq Postgrad Med J.* 2012;11(1):130-5.
 6. Türel I, Özbek H, Erten R, Öner AC, Cengiz N, Yilmaz O. Hepatoprotective and anti-inflammatory activities of *Plantago major L.* *Indian J Pharmacol.* 2009;41(3):120.
 7. Kobeasy I, Abdel-Fatah M, El-Salam SMA, Mohamed ZE-OM. Biochemical studies on *Plantago major L.* and *Cyamopsis tetragonoloba L.* *Int J Biodivers Conserv.* 2011;3(3):83-91.
 8. Cogo LL, Monteiro CLB, Miguel MD, Miguel OG, Cunico MM, Ribeiro ML, et al. Anti-*Helicobacter pylori* activity of plant extracts traditionally used for the treatment of gastrointestinal disorders. *Braz J Microbiol.* 2010;41(2):304-9.
 9. Gomez-Flores R, Calderon C, Scheibel L, Tamez-Guerra P, Rodriguez-Padilla C, Tamez-Guerra R, et al. Immunoenhancing properties of *Plantago major* leaf extract. *Phytother Res.* 2000;14(8):617-22.
 10. Adom MB, Taher M, Mutalabisin MF, Amri MS, Kudus MBA, Sulaiman MWAW, et al. Chemical constituents and medical benefits of *Plantago major*. *Biomed Pharmacother.* 2017;96:348-60.
 11. Tarvainen M, Suomela J-P, Kallio H, Yang B. Triterpene acids in *Plantago major*: Identification, quantification and comparison of different extraction methods. *Chromatographia.* 2010;71(3-4):279-84.
 12. Liu J. Pharmacology of oleanolic acid and ursolic acid. *J Ethnopharmacol.* 1995;49(2):57-68.
 13. Guil-Guerrero JL, Rodríguez-García I. Lipids classes, fatty acids and carotenes of the leaves of six edible wild plants. *Eur Food Res Technol.* 1999;209(5):313-6.
 14. Samuelsen AB, Paulsen BS, Wold JK, Otsuka H, Yamada H, Espevik T. Isolation and partial characterization of biologically active polysaccharides from *Plantago major L.* *Phytother Res.* 1995;9(3):211-8.
 15. Verma S, Singh S. Current and future status of herbal medicines. *Vet World.* 2008;1(11):347-50.
 16. Farokhi F, Khaneshi F. Histopathologic changes of lung in asthmatic male rats treated with hydro-alcoholic extract of *Plantago major* and theophylline. *Avicenna J Phytomed.* 2013;3(2):143.
 17. Saadat S, Boskabadi J, Boskabady MH. Contribution of potassium channels, beta-2-adrenergic and histamine H1 receptors in the relaxant effect of baicalein on rat tracheal smooth muscle. *Iran J Basic Med Sci.* 2019;22(11):1347-52.
 18. Ozaslan M, Didem Karagöz I, Kalender ME, Kilic IH, Sari I, Karagöz A. In vivo antitumoral effect of *Plantago major L.* extract on Balb/C mouse with Ehrlich ascites tumor. *Am J Chinese Med.* 2007;35(05):841-51.
 19. Memarzia A, Amin F, Saadat S, Jalali M, Ghasemi Z, Boskabady MH. The contribution of beta-2 adrenergic, muscarinic and histamine (H1) receptors, calcium and potassium channels and cyclooxygenase pathway in the relaxant effect of *Allium cepa L.* on the tracheal smooth muscle. *J Ethnopharmacol.* 2019;241:112012.
 20. Linden A, Bergendal A, Ullman A, Skoogh B-E, Löfdahl C-G. Salmeterol, formoterol, and salbutamol in the isolated guinea pig trachea: differences in maximum relaxant effect and potency but not in functional antagonism. *Thorax.* 1993;48(5):547-53.
 21. Popa VT, Somani P, Simon V. The effect of inhaled verapamil on resting bronchial tone and airway contractions induced by histamine and acetylcholine in normal and asthmatic subjects 1-4. *Am Rev Respir Dis.* 1984;130(6):1006-13.
 22. Miyahara Y, Kizawa Y, Sano M, Murakami H. Effects of organic and inorganic Ca²⁺-antagonists on acetylcholine-induced contraction in molluscan (*Mytilus edulis*) smooth muscle. *Gen Pharmacol.* 1993;24(6):1419-23.
 23. Buckle D, Arch J, Bowring N, Foster K, Taylor J, Taylor S, et al. Relaxant effects of the potassium channel activators BRL 38227 and pinacidil on guinea-pig and human airway smooth muscle, and blockade of their effects by glibenclamide and BRL 31660. *Pulm Pharmacol.* 1993;6(1):77-86.
 24. Lronards B, Rampart M, Herman A. Selective M3 muscarinic receptor inhibit smooth muscle contraction in rabbit trachea without increasing the release of acetylcholine. *J Pharmacol Exp Ther.* 1992;263:770-3.
 25. Boskabady MH, Boroushaki M, Aslani MR. Relaxant effect of *Portulaca oleraceae* on guinea pig tracheal chains and its possible mechanism (s) of action. *Med Hypotheses Res.* 2004;1:139-47.
 26. Danser A, Tom B, De Vries R, Saxena PR. L-NAME-resistant bradykinin-induced relaxation in porcine coronary arteries is NO-dependent: effect of ACE inhibition. *Br J Pharmacol.* 2000;131(2):195-202.
 27. Shimizu K, Yoshihara E, Takahashi M, Gotoh K, Orita S, Urakawa N, et al. Mechanism of relaxant response to papaverine on the smooth muscle of non-pregnant rat uterus. *J Smooth Muscle Res.* 2000;36(3):83-91.
 28. Slattery MM, Friel AM, Healy DG, Morrison JJ. Uterine relaxant effects of cyclooxygenase-2 inhibitors in vitro. *Obstet Gynecol.* 2001;98(4):563-9.

29. Satoh K, Yamada H, Taira N. Differential antagonism by glibenclamide of the relaxant effects of cromakalim, pinacidil and nicorandil on canine large coronary arteries. *Naunyn Schmiedebergs Arch Pharmacol.* 1991;343(1):76-82.
30. Semenov I, Brenner R. Voltage effects on muscarinic acetylcholine receptor-mediated contractions of airway smooth muscle. *Physiol Rep.* 2018;6(17):13856.
31. Fukunaga K, Kume H, Oguma T, Shigemori W, Tohda Y, Ogawa E, et al. Involvement of Ca²⁺ signaling in the synergistic effects between muscarinic receptor antagonists and β 2-adrenoceptor agonists in airway smooth muscle. *Int J Mol Sci.* 2016;17(9):1590.
32. Oenema TA, Kolahian S, Nanninga JE, Rieks D, Hiemstra PS, Zuyderduyn S, et al. Pro-inflammatory mechanisms of muscarinic receptor stimulation in airway smooth muscle. *Respir Res.* 2010;11(1):130.
33. Gosens R, Rieks D, Meurs H, Ninaber DK, Rabe KF, Nanninga J, et al. Muscarinic M3 receptor stimulation increases cigarette smoke-induced IL-8 secretion by human airway smooth muscle cells. *Eur Respir J.* 2009;34(6):1436-43.
34. Thangam EB, Jemima EA, Singh H, Bag MS, Khan M, Mathias C, et al. The role of histamine and histamine receptors in mast cell-mediated allergy and inflammation: the hunt for new therapeutic targets. *Front Immunol.* 2018;9:1873.
35. Yazdian MA, Khodadoost M, Gheisari M, Kamalinejad M, Ehsani AH. A hypothesis on the possible potential of *Plantago major* in the treatment of urticaria. *Galen Med J.* 2014;3(2):123-6.
36. Stenholm Å, Göransson U, Bohlin L. Bioassay-guided supercritical fluid extraction of cyclooxygenase-2 inhibiting substances in *Plantago major* L. *Phytochem Analysis.* 2013;24(2):176-83.
37. Vogelzang S, Prins H. Patch clamp analysis of the dominant plasma membrane K⁺ channel in root cell protoplasts of *Plantago media* L. Its significance for the P and K state. *J Membr Biol.* 1994;141(2):113-22.